

# Aotearoa New Zealand STI Management Guidelines for Use in Primary Care

## Women who have sex with women

---

### Overview

- Current evidence indicates that women who have sex with women have the same rate of sexually transmitted infections (STIs) as heterosexual women, however the prevalence of particular STIs differs between these groups. Women who have sex with women should not be presumed to be at low or no risk for STIs based on sexual orientation
- Few data are available on the risk for STIs transmitted by sex between women, but risk probably varies by the specific STI and sexual practice (e.g. oral-genital sex; vaginal or anal sex using hands, fingers or penetrative sex toys; and oral-anal sex)
- Women who have sex with women are at risk for acquiring bacterial, viral and protozoal infections from current and former partners, both male and female. Up to 90% of this group of women have had or continue to have male sexual partners
- Prevalence of bacterial vaginosis (BV) is significantly higher for women who have sex with women (20-50%) than exclusively heterosexual women

### Testing advice

As per [sexual health check guideline](#)

- Women who have sex with women are a diverse group with variations in sexual identity, sexual behaviours, sexual practices and risk behaviours. Testing of women who have sex with women will depend on the history,

the clinical picture and risk factors identified

- Routine cervical cancer screening should be offered to all women who have sex with women
- Women who have sex with women should be offered the human papillomavirus (HPV) vaccine in accordance with current guidelines

### **Special considerations:**

- Women who have sex with women frequently have concordant vaginal microbiota with regular female partners, thought to be due to sexual behaviours which transfer vaginal secretions between them. It would be reasonable to suggest that regular female partners of women with BV are tested for BV, and treated if positive, although there is no evidence that this will reduce recurrences

### **Follow up**

If test results are positive, refer to relevant STI management guideline

Even if all test results are negative, use the opportunity to:

- Educate about safer sex practices, and risk minimisation
- Discuss and activate reminders for regular testing according to risk, especially if behaviours indicate the need for more frequent testing
- Discuss need for cervical screening

### **Auditable outcomes**

For 100% of women who have sex with women, testing is discussed openly and opportunities for further engagement are made available

**Endorsement:** These guidelines have been endorsed by the Blood Borne Viruses and Sexually Transmitted Infections Standing Committee (BBVSS).

**Developed by:** the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine (ASHM) ABN 48 264 545 457 | CFN 17788

**Funded by:** The Australian Government Department of Health

**Disclaimer:** Whilst the Australian Department of Health provides financial

assistance to ASHM, the material contained in this resource produced by ASHM should not be taken to represent the views of the Australian Department of Health. The content of this resource is the sole responsibility of ASHM. [www.ashm.org.au](http://www.ashm.org.au)