

Aotearoa New Zealand STI Management Guidelines for Use in Primary Care

Young people

Overview

- Rates of chlamydia and gonorrhoea diagnoses in Aotearoa New Zealand are highest among people aged 15-29 years. Māori and Pacific Peoples carry a significantly higher burden in terms of rates of disease
- Young people are often unaware of their risk of infection and that sexually transmitted infections (STIs) are often asymptomatic
- Negotiating to see all young people on their own, discussing limits of confidentiality at the start of each consultation, minimising costs and routinely offering STI screening (using self-collected samples, when appropriate) to all young people can help overcome barriers to STI testing
- Providing sexual health care to adolescents, particularly younger adolescents, should involve a complete psychosocial (HEADSS) risks and strengths assessment, including screening for any past history of sexual abuse or assault, and an assessment of the young person's capacity for decision making and consent while being aware of child protection issues and mandatory reporting requirements
- The Aotearoa New Zealand Youth'19 survey found that 1% of high school students identified as transgender or non-binary, with a further 0.6% unsure about their gender. These students had compromised health and wellbeing compared with their cisgender peers. Opening discussion about gender identity may begin to address the challenges these young people face, and enable them to access appropriate supports. It is helpful to introduce yourself with your name and the pronouns you use

- It is important to maximise consultations with young people and use them as an opportunity to educate young people about good sexual health practices including safety, condom use, regular screening and contraception

Testing advice

As per sexual health check guideline

- Opportunistically offer STI screening using self-collected swabs (unless symptomatic or examination requested) to all young people at least annually

Special considerations

Privacy

- Ask the young person about the best way to contact them with test results and reminders

Consent

- People aged over 16 years are considered legal adults
- Minors may provide legally effective consent for medical treatment if they are mature enough to understand what is proposed and are capable of expressing their own wishes¹
- A female of any age has the right to consent or refuse to give consent to any medical or surgical procedure for the purpose of terminating her pregnancy²

Child protection

- There is no law in Aotearoa New Zealand that makes reporting of sexual abuse mandatory
- The Children, Young Persons and their Families Act 1989 states that '[the] welfare and interests of the child or young person shall be the first and paramount consideration.'
- The Crimes Amendment Act 2011, Section 195A *Failure to protect child or vulnerable adult* states 'A person is liable for the offence of failing to take

reasonable steps to protect a child or vulnerable adult' if they were aware the person was being abused and failed to take reasonable steps to protect them

- Most health organisations have policies for reporting abuse. Best practice recommends that staff who identify or suspect abuse report their concerns to a statutory agency, the police or Oranga Tamariki
- Decisions to report can be complex and discussion with experienced colleagues and local agencies is often helpful

Follow up

If test results are positive, refer to relevant STI management guideline

- Consider adherence to treatment when deciding on management
- Offer single dose medication where possible

Even if all test results are negative, use the opportunity to:

- Educate about negotiating safer sex, how to use condoms and risk minimisation
- When appropriate, address contraceptive needs
- Vaccinate for hepatitis B and human papillomavirus (HPV) as required
- Offer reminders for regular screening tests
- Discuss need for cervical screening if aged over 25 years

Auditable outcomes

- 90% of young people attending the practice are tested according to these guidelines
- 100% of young people tested have hepatitis B and HPV vaccination status recorded

Useful patient resources

[Just the facts](#)

References

1. Morris KA, editor. Coles Medical Practice in New Zealand 13th edition 2017.
2. Medical Council of New Zealand (MCNZ). Appendix 1. Information, choice of treatment and informed consent. February 2019.

Endorsement: These guidelines have been endorsed by the Blood Borne Viruses and Sexually Transmitted Infections Standing Committee (BBVSS).

Developed by: the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine (ASHM) ABN 48 264 545 457 | CFN 17788

Funded by: The Australian Government Department of Health

Disclaimer: Whilst the Australian Department of Health provides financial assistance to ASHM, the material contained in this resource produced by ASHM should not be taken to represent the views of the Australian Department of Health. The content of this resource is the sole responsibility of ASHM. www.ashm.org.au