

# Aotearoa New Zealand STI Management Guidelines for Use in Primary Care

## Vaginal discharge

---

### Overview

- Vaginal discharge is a common complaint
- Vaginal discharge consists of cervical mucous, vaginal transudate, vaginal epithelial cells and vaginal flora
- The most common cause of vaginal discharge in people of reproductive age is normal physiological discharge
- Consider other causes with history, examination and investigations

### Cause

Bacterial vaginosis

Vulvovaginal candidiasis

### Sexually transmitted infections

- *Chlamydia trachomatis*
- *Neisseria gonorrhoeae*
- *Trichomonas vaginalis*
- Herpes simplex virus (HSV) (cervical infection)

### Non-infectious causes

- Hormonal contraception, physiological, cervical ectropion and cervical polyps
- Foreign body (e.g. retained tampon)

- Atrophic vaginitis in lactating and postmenopausal women or women on hormonal contraception e.g. hormone releasing intrauterine device (IUD), Mirena, Depo-Provera or more rarely combined oral contraception
- Desquamative inflammatory vaginitis (DIV) or aerobic vaginitis

## **Rare**

- Bacterial vaginitis e.g. streptococcal or staphylococcal
- Cervical malignancy

## **Clinical presentation**

### **Symptoms**

- Volume of discharge
- Change in characteristics e.g. yellow, green or white, curd-like, frothy
- Odour – fishy smelling, malodorous
- Associated vulval symptoms – itching, tenderness, rash
- Other relevant symptoms – abnormal vaginal bleeding, dyspareunia, pelvic pain may indicate pelvic inflammatory disease (PID)

### **Indications for testing**

- Change from usual vaginal discharge

### **Recommended tests**

- Examination recommended, including inspection of external genitalia, speculum examination of cervix and vagina, and bimanual palpation (if indicated)
- Specifically, examine for:
  - Characteristics of discharge (colour, consistency, distribution, volume and odour)
  - Cervicitis
  - Vaginitis
  - Vulvitis
  - Ulceration

- Signs of PID
- foreign body (e.g. tampon or condom)

## Investigations

- As per sexual health check guideline

## Management

- Syndromic management of vaginal discharge described below
- Refer to appropriate guideline if results known.

Examination findings	Treatment
Vulvovaginitis Thick white curd-like discharge Itching and tenderness	<ul style="list-style-type: none"> <li>· Presumptive vulvovaginal <u>candidiasis</u></li> <li>· Clotrimazole 2% vaginal cream 3-day course</li> <li>· OR clotrimazole 1% vaginal cream 6 nights (preferred regimen for pregnancy as less likely to respond to shorter course)</li> <li>· OR fluconazole 150 mg orally, as a single dose (not in pregnancy)</li> </ul>
Fishy smelling white or grey adherent discharge NO vulvovaginitis	<ul style="list-style-type: none"> <li>· Presumptive <u>bacterial vaginosis</u></li> <li>· Metronidazole 400 mg orally, twice daily for 7 days (preferred regimen for pregnancy)</li> <li>· OR metronidazole 2 g orally, as a single dose if adherence an issue (less effective)</li> </ul>
Offensive purulent frothy discharge +/- vulvovaginitis	<ul style="list-style-type: none"> <li>· Presumptive <u>trichomoniasis</u></li> <li>· Metronidazole 400 mg orally, twice daily for 7 days (preferred regimen for pregnancy)</li> <li>· OR metronidazole 2 g orally, as a single dose</li> </ul>

## Special situations

Situation	Recommendation
Foreign body/retained tampon	Remove foreign body Metronidazole 400 mg orally, twice daily for 7 days May cause a discharge similar to bacterial vaginosis, and there may often be cervicitis if present for some time

Peri or postmenopausal women	<p>Consider vaginal oestrogen cream for symptomatic women with negative tests</p> <p>Note: <u>Bacterial vaginosis</u> may be reported in this setting due to the altered microbiome of menopause, and topical oestrogen may be used as treatment</p>
------------------------------	--

- Treat initially empirically as above, and then based on results when they become available
- More than one pathology may be present e.g. bacterial vaginosis and vulvovaginal candidiasis
- If sexually transmitted infection (STI) suspected, advise no unprotected sexual intercourse until results available
- Vaginal creams can damage latex condoms

### **Partner notification and management of sexual contacts**

- Contact tracing is only required for STIs – see appropriate guideline

### **Follow up**

<ul style="list-style-type: none"> <li>• Not required if tests negative, and symptoms resolve</li> <li>• If test positive, refer to appropriate guideline</li> </ul>
<p style="text-align: center;"><b>Test of cure</b></p> <ul style="list-style-type: none"> <li>• Refer to appropriate guideline if test positive</li> </ul>
<p style="text-align: center;"><b>Retesting</b></p> <ul style="list-style-type: none"> <li>• Refer to appropriate guideline if test positive</li> </ul>

### **Indications for specialist referral**

**Referral to or discussion with a sexual health specialist is recommended for:**

- Persistent or recurrent vaginal discharge
- Lack of response to appropriate treatment
- Complicated clinical situations for further management

### **Auditable outcomes**

## **Useful patient resources**

What's going on down there

**Endorsement:** These guidelines have been endorsed by the Blood Borne Viruses and Sexually Transmitted Infections Standing Committee (BBVSS).

**Developed by:** the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine (ASHM) ABN 48 264 545 457 | CFN 17788

**Funded by:** The Australian Government Department of Health

**Disclaimer:** Whilst the Australian Department of Health provides financial assistance to ASHM, the material contained in this resource produced by ASHM should not be taken to represent the views of the Australian Department of Health. The content of this resource is the sole responsibility of ASHM. [www.ashm.org.au](http://www.ashm.org.au)